

Earth-mass dark halos are torn into dark mini-streams by stars

HongSheng Zhao¹, *James Taylor², †Joseph Silk³ & Dan Hooper³ ‡

Communications Arising from *Letter to Nature* by Diemand et al.

The promising neutralino dark matter particles generically condense into numerous earth-mass dark halos with smaller substructures suppressed by free streaming[1]. The recent *Letter to Nature*[2] claims that these 0.01pc-sized dense halos emerged at redshifts 60–26 are rarely destructed inside galaxies, hence the nearest halos at about 0.1pc from Earth are bright in annihilation-powered gamma-rays, but are inaccessible to direct detections. However, most mini-halos reaching solar neighbourhood should experience strong impulses by individual stars in the Galactic disk and bulge, and have been torn into pc-long tidal streams over a Hubble time with only modest overdensity, reducing indirect annihilation signals. Sweeping across the solar system

*¹PPARC Advanced Fellowship at University of St Andrew's, KY16 9SS, UK. Young Overseas Scholar at Beijing Observatory. E-mail: hz4@st-and.ac.uk

†²Astronomy Department, Caltech, mail code 105-24, 1200 East California, Pasadena CA 91125, Email: jet@astro.caltech.edu

‡³Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH Email: silk@astro.ox.ac.uk, hooper@astro.ox.ac.uk

per century, mini-streams leave directional and temporal signatures for direct searches of neutralinos.

When a minihalo of a mean density $\rho \equiv \frac{3m(r)}{4\pi r^3}$ inside radius r encounters a star of mass $M_* \gg m(r)$ with a high relative velocity V_r and impact parameter $b > r$, the minihalo is stripped outside a radius r if the specific heating $\frac{(\delta v)^2}{2} \times \frac{2}{3}$ (a projection factor of the differential impulse δv perpendicular to and along the star's almost straight path)[3] is greater than the specific binding energy $\frac{Gm(r)}{2r}$ at radius r , where δv is the tidal force $\frac{GM_*r}{b^3}$ times the effective duration of the impulse $\frac{2b}{V_r}$. Hence only the dense part of the minihalo survive with $G\rho > k \left(\frac{GM_*}{S}\right)^2$ where the factor $k = 2\pi$, and $S \equiv (\pi b^2)V_r$ is the volume plunged through by the minihalo per unit time. We estimate $S \leq H_0 \langle n_* \rangle$ by requiring fewer than one disruptive encounter per Hubble time H_0^{-1} in a sea of normal stars of mass M_* with a time and space averaged star density $\langle n_* \rangle M_*$, which is estimated as[5] either $\sim 0.001 M_\odot \text{ pc}^{-3} \left(\frac{16 \text{ kpc}}{2H}\right)$ or $\sim 0.012 M_\odot \text{ pc}^{-3} \left(\frac{16 \text{ kpc}}{2H}\right) \left(\frac{8 \text{ kpc}}{R}\right)^2$ by averaging either an old stellar local disk density $\sim 20 M_\odot \text{ pc}^{-2}$ over a column of orbital height $2H$, or an old stellar disk and bulge mass $4 \times 10^{10} M_\odot$ over a cylinder of orbital radius R and height $2H$. The survival minihalos are regions of overdensity above $\frac{\rho_{\text{surv}}}{\rho_{\text{crit}}(z=60)} = 200 \left[\frac{\langle n_* \rangle M_*}{0.001 M_\odot \text{ pc}^{-3}} \right]^2$ where we take the prefactor $k \sim 10$ ([4] found $k \sim 30$ in binary disruption simulations including head-on encounters and a series of gradual heating with large impact parameters). Scaled to the background density ρ_{crit} at the emerging redshift $z = 60$, the $10^{-6} M_\odot$ minihalos proposed by[2] have low overdensities ~ 4 and 200 inside half-mass radius 0.01 pc and their 0.001 pc resolution limit (enclosing of order 5 percent of the minihalo mass, cf.

their Fig.2) respectively, hence these minihalos should be mostly stripped by encounters long ago. The resupply of minihalos is insignificant locally since the local overdensity collapsed mainly at high redshift.

More sophisticated semi-analytical models of the massloss[6] also predicts that mini-halos on polar orbits would lose a few percent of the mass per orbit if they spend a few percent of the orbital period in strong tide regime near disk stars, hence over a Hubble time ($\sim 20\text{--}70$ orbits at the solar radius), a minihalo with 1m/s escape velocity is stripped nearly completely and stretched hundred fold to a modest overdensity region of several pc long. These dark streams roughly fills the local volume. The Earth is likely crossing one of these 0.02pc wide mini-streams on time scales of centuries. Sensitive, long-term direct detection experiments might expect to see slow variations of order 1% per year. Finally, the fact that local minihalos on planar orbits with small pericentres are preferentially destructed could imprint an interesting directional or phase modulation on direct detection signals. While the smoothing of the local halo increases the chance of direct detection, it simultaneously reduces the indirect signals of annihilation [7], which depend on neutralino density squared, hence is proportional to the fraction of dark substructures in the local density and their contrast to the smooth component. The boost factor drops from $\sim 15 - 45$ for unstripped concentrated substructures to $\sim 4 - 12$ if most minihalos are disrupted and stretched, although this boost is still larger than previous estimates [1].

References

- [1] Berezhinsky, V., Dokuchaev, V. and Eroshenko, Y. Small-scale clumps in the galactic halo and dark matter annihilation. *Phys. Rev. D* **68**, 103003 (2003).
- [2] Diemand, J., Moore, B. and Stadel, J. Earth-mass dark-matter haloes as the first structures in the early Universe. *Nature* **433**, 389 (2005).
- [3] Binney J.J. & Tremaine S. 1987, *Galactic Dynamics*, Princeton University Press.
- [4] Bahcall J., Hut P., Tremaine S. The maximum mass of the objects that constitute the unseen disk material. *Astrophys. J.* **290**, 15 (1985).
- [5] Robin A.C., Reyl   C., Derrier   S., Picaud S., A synthetic view on structure and evolution of the Milky Way. *Astronomy & Astrophys.* **409**, 523 (2003).
- [6] Taylor, J. E. & Babul, A. The dynamics of sinking satellites around disk galaxies: a poor man's alternative to high-resolution numerical simulations. *Astrophys. J.* **559**, 716 (2001).
- [7] J. E. Taylor and J. Silk, The clumpiness of cold dark matter: Implications for the annihilation signal, *Mon. Not. Roy. Astron. Soc.* **339**, 505 (2003).